

## Dietary enrichment of broiler chicken with omega-3 fatty acids and beneficial role in human cardiovascular health: A Review

M.S.M.Nafees and M.Pagthinathan

Department of Animal Science, Faculty of Agriculture, Eastern University,  
Chenkalady, Sri Lanka  
mohamednafeez@yahoo.com

### ABSTRACT

Broiler meat consumption in the world increases annually with the development of modern diet pattern. Saturated fatty acid content of the meat causes high risk of cardiovascular disease in human. Fish is an important source of omega-3 fatty acids, which reduce the risk of cardiovascular disease in human. However, access to food fish, which are rich in omega-3 fatty acids, is questionable for many people in the world. Fortunately, studies revealed that the dietary enrichment of broiler chicken ration with omega-3 fatty acids enriched the broiler chicken meat with omega-3 fatty acids. Hence, consumption of such broiler chicken meat may supplement the scarcity of consumption of omega-3 fatty acids rich foods viz. fish. Moreover, dietary enrichment of omega-3 fatty acids improves carcass by reducing the abdominal fat deposition in broilers. However, shelf-life of broiler chicken meat that is enriched with omega-3 fatty acids low due to high levels of lipid oxidation. However, usage of antioxidants improves the shelf-life of omega-3 fatty acid enriched broiler chicken meat.

### Introduction

Animal source foods are complete and nutrient-dense, and provide high quality protein and bio-available micronutrients viz. iron, zinc vitamin A, vitamin B<sub>12</sub> and calcium, particularly for children and pregnant and lactating mothers (FAO, 2011). In particular, consumption of meats prevents anaemia and enhances iron absorption from plant based foods (Walker *et al.*, 2005). Hence, different communities in the world regard meat as a desirable food with high nutritive value (FAO, 1992;

Michael and Bambrick, 2005). Irrespective of cultural taboos for the consumption of meats, poultry meat is accepted in many cultures (FAO, 2011), particularly broiler chicken meat (Chashnidel *et al.*, 2010). In developing countries, a great rise in the production of poultry has been observed (Fig. 1) and expected to continue in future (Speedy, 2003).




Fig 1. Growth in production of agricultural products in developing countries (1975–2005)  
Source: FAO (2008).

Compare to pigmeat, bovine meat, and ovine meat, world poultry meat production has grown at a higher rate for last ten years (Fig. 2), and it is expected to rise by 1.5% in 2015 (FAO, 2015). According to FAO (2011), growth in world poultry meat consumption will be 225% by 2050 (Fig. 3). In particular, developing countries have shown a higher growth in broiler production (Chang, 2007). Globally, poultry meat contributes around 35% of all meats (FAO, 2015), whereas in Sri Lanka 75% of the per capita meat consumption is contributed by poultry meat (Department of Animal Production and Health, 2011), mainly by the chicken (De Silva *et al.*, 2010).




Fig. 2 World meat production trends in past ten years  
Source: FAO Food Outlook (2006-2015).




Fig 3. Projected growth in world meat consumption  
Source: FAO (2011).

On the contrary, meats are the primary source of saturated fats, which are responsible for high risk of cardiovascular disease, diabetes mellitus and cancers in human (McMichael and Bambrick, 2005). Moreover, modern agribusiness and food processing industries change the nutrient content and structure of many food items. For example, hydrogenation of foods changes the double bonds of fatty acid from *cis* to *trans* and result with increased *trans* fatty acids in processed foods. This structural change increases the risk of coronary heart disease (CHD) in human (Simopoulos, 1999). Furthermore, modern diets have high saturated fatty acids (SFAs) and low mono-unsaturated fatty acids (MUFAs) and poly-unsaturated fatty acids (PUFAs) (de Witt *et al.*, 2009). Myristic and palmitic acids are the principal dietary SFAs that increase the blood cholesterol level (Bender, 1992). Arterial plaques, which lead to partial blockage of the blood vessels, are caused by oxidized (rancid) form of cholesterol (Farrell, 2010).

In these contexts, there are growing interests in the world in modifying the cholesterol content and fatty acid compositions of poultry products (Hargis and Van Elswyk, 1993; Sacks, 2002; Salma *et al.*, 2007) to produce superior health quality food. Recent researches also deal with reducing fat, cholesterol, and SFA contents of poultry meat by dietary supplementation of garlic, copper,  $\alpha$ -tocopherol acetate and omega-3 fatty acid (Salma *et al.*, 2007). The latter has utmost importance due to its human health benefits. However, consumption of omega-3 fatty acid by human is rare. Hence, there are several studies have been carried out in the world to enrich the favourite human foods *viz.* poultry meat with omega-3 fatty acids so as to provide health benefits to the human.

This article is intended to review the studies related to dietary enrichment of broiler chicken with omega-3 fatty acids and their impacts on productive parameters of broiler chicken and health condition of broiler chicken meat consumers.

### **Omega-3 fatty acids**

Dietary fat consists of both SFA and unsaturated fatty acids (UFA). The latter further divided into MUFA and PUFA. Based on the chemical nature, PUFAs are further classified into omega-3 fatty acids (also called as  $\omega$ -3 fatty acids or n-3 fatty acids) and omega-6 fatty acids (Din *et al.*, 2004). Thus, omega-3 fatty acids are one of two families of essential PUFAs. The term omega-3 (also  $\omega$ -3 or n-3) indicates the first double bond exists as the third carbon-carbon bond from the terminal methyl end of the carbon chain. Eicosapentaenoic acid (EPA), docosahexaenoic acid (DHA) and alpha-linolenic acid (ALA) are the major omega-3 fatty acids useful for humans. EPA (C20:5 n-3) and DHA (C22:6 n-3) are rich in marine sources *viz.* fish and shellfish while ALA (C18:3 n-3) is rich in plant sources *viz.* flaxseed (linseed), soybean, walnut, canola and rapeseed oils (Stone, 1996; Mellor, 2005).

### **Importance of omega-3 fatty acids in human health**

Consumption of long chain omega-3 fatty acids *viz.* EPA and DHA reduces risk of cardiovascular diseases (CVD) in human by lowering the blood pressure and heart beat rate (Kromhout *et al.*, 1995; Connor, 2000; Kris-Etherton *et al.*, 2002; AHRQ, 2004; Din *et al.*, 2004; Grashorn, 2005; Mellor, 2005; Hooper *et al.*, 2006; Schacky and Harris, 2007; Betti *et al.*, 2009; Mirghelenj *et al.*, 2009). Moreover, the long chain omega-3 fatty acids reduce serum triglycerides, thrombotic tendency, inflammation, and arrhythmias and improves endothelial function, insulin sensitivity, paraoxonase concentrations, and plaque stability in human (Geelen *et al.*, 2004).

Nevertheless, true essential fatty acids (EFAs) *viz.* linoleic acid (C18:2 n-6) and ALA (Mellor, 2005) should be converted into EPA and DHA to be effective in human body (Gerster, 1998; Brenna, 2002; AHRQ, 2004). However, the rate of conversion is only around 2% (Mellor, 2005) to 5% (Brenna, 2002) as humans lack the necessary enzymes to convert omega-6 fatty acids to omega-3 fatty acids (Din *et al.*, 2004). Hence, it is very important to have dietary EPA and DHA through meat based diet such as poultry meat, marine and freshwater fish, *etc.* to get advantages of omega-3 PUFAs.

### **Fish as a rich dietary source of EPA and DHA**

Vegetable oils are rich in ALA, whereas fish and fish oils are rich in EPA and DHA (Sargent, 1997; Din *et al.*, 2004; Mirghelenj *et al.*, 2009). However, amount of EPA and DHA varies among fish species (Table 1) and environmental factors. WHO and NATO recommended 0.3 to 0.5 g/d of EPA+DHA and 0.8 to 1.1 g/d of ALA as a dietary inclusion level of omega-3 fatty acids for a healthy person (Kris-Etherton *et al.*, 2002). Furthermore, American Heart Association (AHA) recommended 1 g/d of EPA+DHA for CHD patients (Kris-Etherton *et al.*, 2002; Schacky and Harris, 2007). Although this level of EPA and DHA could be obtained through fish consumption, the required intake may be difficult to achieve and sustain in long term (Kris-Etherton *et al.*, 2002) due to seasonal availability, affordability and consumer preference of fish (Hargis and Van Elswyk 1993). Hence, supply of these PUFAs is easier if they are incorporated with widely consumed food items *viz.* meats. In this context, broiler chicken meat is ideal for the enrichment with essential fatty acids.

In poultry, lipid digestion takes place in the small intestine, where the pancreatic lipase breakdowns the triacylglycerols into 2-monoacylglycerols and free fatty acids (Nieto and Ros, 2012). These products are formed into micelles when linked to the conjugated bile salts. Monoglycerides and long-chain unsaturated fatty acids have higher ability to form the micelles with bile salts than saturated fatty acids due to their characteristic low polarity (Baião and Lara, 2005). Because of pancreatic lipase, the mixed micelles are absorbed without any alteration in the composition of fatty acids. Hence, the dietary lipid sources have a direct effect on the fatty acid composition of poultry products (Baião and Lara, 2005; Nieto and Ros, 2012).

As the tissues of poultry could simply be enriched with unsaturated fatty acids by increasing their proportion in the diet (Nieto and Ros, 2012), enrichment of broiler chicken meat with omega-3 fatty acids could be an alternative source for fish (Hargis and Van Elswyk 1993).

**Table 1. Long chain omega-3 fatty acid contents of selected fin-fish and shellfish species**

	EPA+DHA content (g/100g of edible portion of fish)	Fish required to provide 1 g of EPA+DHA(g)
<b>Fin-fish</b>		
Catfish	0.18	556
Cod	0.28	357
Tuna (fresh)	0.28 - 1.51	66 - 357
Tuna (canned)	0.31	323
Mackerel	0.4 - 1.85	54 - 250
Halibut	0.47 - 1.18	85 - 213
Rainbow trout	1.15	87
Sardines	1.15 - 2	50 - 87
Atlantic salmon	1.28 - 2.15	42.5 - 70.9
Atlantic herring	2.01	50
<b>Shellfish</b>		
Scallop	0.2	500
Shrimp	0.32	313
Oyster	0.44	227

Source: Din *et al.* (2004).

### Dietary enrichment of broiler chicken meat with omega-3 fatty acids

Chickens modify their lipid profile within a week of feeding a new diet enriched with lipid sources (Lopez-Ferrer *et al.*, 2001). Dietary fatty acids could be absorbed and deposited in the body tissue of broiler chickens without any modifications. Thus, lipid composition of broiler chicken meat could be altered according to the fatty acid profile of the dietary lipid sources (de Witt *et al.*, 2009; Hugo *et al.*, 2009b). This ensures the potential to increase the consumption of omega-3 PUFAs through broiler chicken meat (Coetzee and Hoffman, 2002).

Unfortunately, high growth rate of modern broiler chicken breeds has been associated with increased fat deposition. Lipids in broiler chicken are mainly deposited as abdominal fat (20%) and subcutaneous fat (18%), and skeleton fat (15%), fats on liver and feather (2.5%), and fats on carcass (40%) as intermuscular fat, intramuscular fat and subcutaneous fat. Abdominal and subcutaneous fats are the main sources of waste in the slaughterhouse (Tumová and Teimouri, 2010). However, a minimum quantity of intramuscular fat is necessary for an optimal sensory quality because of its positive influence on succulence, tenderness and flavour (Guo-Bin *et al.*, 2010; Tumová and Teimouri, 2010; Pegg and Shahidi, 2012).

Dietary enrichment of broiler chicken meat with unsaturated fatty acids positively improved the

intramuscular fat deposition. Baião and Lara (2005) reported that the broiler chicken fed with a diet containing sunflower oil (unsaturated fat) and bovine/swine fat (saturated fat) caused higher accumulation of intramuscular fat and abdominal fat. However, broilers fed with a diet containing 8% sunflower oil had significantly less amount of abdominal fat deposits than those fed with a diet containing 8% beef tallow (saturated fat). Thus, the location of fat deposition in broiler chicken depends on the kind of fatty acid added to the diet.

Past studies revealed that incorporation of omega-3 PUFA rich sources *viz.* fish oil (Lopez-Ferrer *et al.*, 2001; Basmacıođlu *et al.*, 2003; Mirghelenj *et al.*, 2009) into broiler chicken diet enriched the chicken meat with omega-3 PUFAs (Hargis and Van Elswyk, 1993; Leskanich and Noble, 1997; Moghadasian, 2008; Zuidhof *et al.*, 2009; Zduñczyk and Jankowski, 2013). Table 2 shows that different sources of long-chain (LC) *n-3* PUFAs used in the broiler chicken ration have the impact on the fat deposition of the broiler chicken meat (Zduñczyk and Jankowski, 2013).

In addition to fish oil, marine algae (Basmacıođlu *et al.*, 2009) or precursors of omega-3 PUFAs *viz.* ALA through canola oil (Shahriar *et al.*, 2007; Salamatdoustnobar *et al.*, 2008), linseeds, flaxseed or similar oils (Lopez-Ferrer *et al.*, 1999), also used for dietary manipulation of broiler chicken.

**Table 2. Fatty acid composition of diets with different sources of PUFAs and whole-body concentration of PUFAs in chickens**

Fatty acid composition (%)	Sources of LC <i>n-3</i> PUFAs			
	Palm oil	Soybean oil	Linseed	Fish oil
In the diet				
<i>n-6</i> PUFAs	14.7	24.11	14.5	12.4
<i>n-3</i> PUFAs	0.9	2.4	14.1	9.8
<i>n-6/n-3</i>	16.7	10.0	1.0	1.3
In the body of chicken				
<i>n-6</i> PUFAs	11.9	22.7	14.4	10.5
<i>n-3</i> PUFAs	0.7	2.2	13.4	6.9
<i>n-6/n-3</i>	17.0	10.3	1.1	1.5
Accumulation (% of net intake)				
<i>n-6</i> PUFAs	73.1	74.7	85.2	73.0
<i>n-3</i> PUFAs	63.9	64.3	65.3	58.4

Source: Zduńczyk and Jankowski (2013).

#### **Dietary omega-3 PUFA on bodyweight gain of broiler chickens**

Dietary omega-3 PUFA alters bodyweight (BW) of broiler chickens. Studies showed that 1% dietary conjugated LA (CLA) increased bodyweight gain (BWG) of broiler chickens (Lopez-Ferrer *et al.*, 2001; Aydin, 2007). Similarly, dietary omega-3 PUFA supplemented with vitamin E and organic Se improved BW of broiler chickens (Malayođlu *et al.*, 2009). According to Haug *et al.* (2007), a high level of dietary organics increased the concentration of EPA and DHA in the broiler chicken muscle tissues. However, enrichment of omega-3 PUFA may increase energy content of the diet, thereby excess energy could be deposited as fat. In contrast, Salamatdoustnobar *et al.* (2008) and Szymczyk *et al.* (2001) observed that the dietary omega-3 PUFA reduced the abdominal fat content of broiler chickens.

Broiler chicken gets fat deposition by means of exogenous (from the diet) or endogenous (synthesized in the liver) or from skeleton. Absorbed dietary fat is packed into portomicrons and non-esterified fatty acids which pass directly into the hepatic portal blood supply, whereas fat synthesized in the liver is transported to other tissues in the form of triglyceride-rich very low density lipoprotein (Tumová and Teimouri, 2010). Since dietary omega-3 PUFA lowers the low-density lipoprotein (LDL) levels in blood stream of broiler chickens, fat deposition may be

reduced. Because LDL transfers cholesterol from liver to tissues, thereby it increases the tissue fat deposition. Moreover, there is an effect of genotype on the dressing percentage and growth of breast muscle of broiler chickens enriched with omega-3 PUFA (de Witt *et al.* 2009).

#### **Dietary omega-3 PUFA on lipid oxidative stability of broiler chicken meat**

Lipid oxidation in muscle and fat tissues affects the shelf-life of meat and meat products thereby deteriorate quality of meat products. Membrane lipids (phospholipids) in the meat have a greater share of PUFAs, which are most prone to oxidation (Pegg and Shahidi, 2012). However, dietary enrichment of broiler chicken meat with PUFAs has become as a popular practice in the world due to human health concerns. Hence, the chance of lipid oxidation in broiler chicken meat is high. Studies revealed that the lipid oxidative stability is low in omega-3 PUFA enriched broiler chicken (Hugo *et al.*, 2009b) especially, colour, flavor and shelf life of meat are affected (Smet *et al.*, 2008; Betti *et al.*, 2009). Although, dietary fish oil in broiler chicken ration causes "fishy taint" in broiler chicken meat through the oxidation of omega-3 PUFA which in turn reduce the feed intake of birds (Chashnidel *et al.*, 2010) and consumer preference of chicken meat (Hugo *et al.*, 2009a).

More than 2% fish oil in the broiler chicken ration reduces consumer preference of chicken meat fed with that ration due to fishy smell in cooked meat (Mirghelenj *et al.*, 2009). Moreover, oxidation of highly PUFA enriched meat may harm human health (Lopez-Ferrer *et al.*, 2001).

### Improving meat quality

Favourable enrichment of meats with omega-3 PUFAs depends on the dose of fish oil or other omega-3 rich PUFA sources (Leskanich and Noble, 1997; Bou *et al.*, 2004). In this context, chicken diet should be enriched with less than 15-20 g fish oil per kg feed or less than 120 g fish meal per kg feed to minimise unpleasant aromas and flavours (Hargis and Van Elswyk, 1993; Leskanich and Noble, 1997; Howe *et al.* 2002). However, combining fish oils and vegetable oils and/or seeds could favourably increase the omega-3 PUFAs in chicken meat (Bouet *al.* 2005). Moreover, usage of quality fish oil and addition of antioxidants *viz.* vitamin E in the broiler chicken diets could improve oxidative stability of such meat.

### Summary

Biologically important omega-3 PUFAs are EPA, DHA and ALA. The latter is abundant in plant sources while, others are rich in fish. Enrichment of broiler chicken meat with omega-3 PUFA is possible through dietary treatment. Dietary enrichment of omega-3 PUFA improves body weight gain in broiler chickens whereas consumption of omega-3 PUFA enriched broiler chicken meat prevents risk of cardiovascular disease in human.

### References

- AHRQ. (2004). Effects of Omega-3 Fatty Acids on Cardiovascular Risk Factors and Intermediate Markers of Cardiovascular Disease, Number 93: Agency for Healthcare Research and Quality, U.S. Department of Health and Human Services, 540 Gaither Road, Rockville.
- Aydin, R. (2007). Effect of dietary oils and conjugated linoleic acid on the growth performance of broilers vaccinated with the La Sota Newcastle vaccine. *South African Journal of Animal Science*, 37:74-80.
- Baião, N.C. and L. J. C. Lara (2005). Oil and Fat in Broiler Nutrition. *Brazilian Journal of Poultry Science*, 7:129-141.
- Basmacıođlu, H., M. Çabuk., K. Ünal., K. Özkan., S. Akkan and H. Yalçın (2003). Effects of dietary fish oil and flaxseed on cholesterol and fatty acid composition of egg yolk and blood parameters of laying hens. *S. Afr. J. Anim. Sci.*, 33: 265-273.
- Basmacıođlu, H., S. Özkan., S. Koçtürk., G. Oktay and M. Ergül (2009). Dietary vitamin E (α-tocopheryl acetate) and organic selenium supplementation: performance and antioxidant status of broilers fed n-3 PUFA-enriched feeds. *South African Journal of Animal Science*, 39(4): 274-285.
- Bender, A. (1992). *Meat and meat products in human nutrition in developing countries, food and nutrition paper 53*: animal production and health division and the food policy and nutrition division, Food and Agriculture Organization of United Nations (FAO), Rome.
- Betti, M., B. L. Schneider., W. V. Wismer., V. L. Carney., M. J. Zuidhof and R. A. Renema (2009). Omega-3-enriched broiler meat: 2. Functional properties, oxidative stability, and consumer acceptance. *Poultry Science*, 88:1085-1095.
- Bou, R., F. Guardiola., A. Tres., A. C. Barroeta and R. Codony (2004). Effect of dietary fish oil, alpha-tocopheryl acetate, and zinc supplementation on the composition and consumer acceptability of chicken meat. *Poultry Science*, 83: 282-292.
- Bou, R., F. Guardiola., A. C. Barroeta and R. Codony (2005). Effect of dietary fat sources and zinc and selenium supplements on the composition and consumer acceptability of chicken meat. *Poultry Science*, 84: 1129-1140.

- Brenna, J.T. (2002). Efficiency of conversion of alpha-linolenic acid to long chain n-3 fatty acids in man. *Curr. Opin. Clin. Nutr. Metab. Care* 5:127–132.
- Chang, H. (2007). Overview of the World Broiler Industry: Implications for the Philippines. *Asian Journal of Agriculture and Development*, 4:67-82.
- Chashnidel, Y., H. Moravej., A. Towhidi., F. Asadi and S. Zeinodini (2010). Influence of different levels of n-3 supplemented (fish oil) diet on performance, carcass quality and fat status in broilers. *African Journal of Biotechnology*, 9:687-691.
- Coetzee, G.J.M. and L. C. Hoffman (2002). Effects of various dietary n-3/n-6 fatty acid ratios on the performance and body composition of broilers. *S. Afr. J. Anim. Sci.*, 29: 175-184.
- Connor, W.E. (2000). Importance of n-3 fatty acids in health and disease. *Am J Clin Nutr*, 71:171S–175S.
- Department of Animal Production and Health. (2011). Key Statistics 2011, P. O. Box 13, Peradeniya, Sri Lanka.
- De Silva, P.H.G. J., N. S. B. M. Atapattu and A. L. Sandika (2010). A Study of the Socio - Cultural Parameters Associated With Meat Purchasing and Consumption Pattern: A Case of Southern Province, Sri Lanka. *The Journal of Agricultural Sciences*, 5(2): 71-79.
- de Witt, F.H., S. P. Els., A. Hugo., H. J. van der Merwe, and Fair, M.D. (2009). Influence of dietary lipid sources on carcass traits of broilers. *South African Journal of Animal Science*, 39:49-52.
- Din, J.N., D. E. Newby and A. D. Flapan (2004). Omega 3 fatty acids and cardiovascular disease—fishing for a natural treatment. *BMJ*, 328:30-35.
- FAO. (1992). *Meat and meat products in human nutrition in developing countries*, Vol. 53: Animal Production and Health Division and Food Policy and Nutrition Division of FAO, Food and Agriculture Organization of United Nations (FAO), Rome, Italy.
- FAO. (2008). *Poultry in the 21st Century: avian influenza and beyond*. Animal Production and Health Proceedings, No. 9: Food and Agriculture Organization of United Nations (FAO), Rome, Italy.
- FAO. (2011). *World Livestock 2011 – Livestock in food security*. Office of Knowledge Exchange, Research and Extension, Food and Agriculture Organization of United Nations (FAO), Rome, Italy.
- FAO. (2015). *Food Outlook 2015 (October 2015)*. Food and Agriculture Organization of United Nations (FAO), Rome, Italy.
- Farrell, D. (2010). The role of poultry in human nutrition: How important is cholesterol in eggs? Poultry Development Review, Food and Agriculture Organization of the United Nations (FAO), Rome.
- Geelen, A., I. A. Brouwer., P. L. Zock and M. B. Katan (2004). Antiarrhythmic effects of n-3 fatty acids: evidence from human studies. *Curr Opin Lipidol*, 15:25-30.
- Gerster, H. (1998). Can adults adequately convert alpha-linolenic acid (18:3n-3) to eicosapentaenoic acid (20:5n-3) and docosahexaenoic acid (22:6n-3). *Int. J. Vitam. Nutr. Res.*, 68:159–173.
- Grashorn, M.A. (2005). Enrichment of eggs and poultry meat with biologically active substances by feed modifications and effects on the final quality of the product. *Pol. J. Food Nutr. Sci.*, 14(55):15–20.
- Guo-Bin, C., L. Li-Li., Z. Xue-Yu., W. Ke-Ilua., C. Rong., L. De-Qin and C. Guo-Hong (2010). Development Rule of Intramuscular Fat Content in Chicken. *Journal of Animal and Veterinary Advances*, 9(2): 297-298.

- Hargis, P.S and M. E. Van Elswyk (1993). Manipulating the fatty acid composition of poultry meat and eggs for the health conscious consumer. *World's Poultry Science Journal*, 49:251-264.
- Haug, A., S. Eich-Greatorex., A. Bernhoft., J. P. Wold., H. Hetland., O. A. Christophersen and Y. Sogn (2007). Effect of dietary selenium and omega-3 fatty acids on muscle composition and quality in broilers. *Lipids in Health and Disease*, 6:29.
- Howe, P., J. Downing., B. Grenyer., E. Grigonis-Deane and W. Bryden (2002). Tuna fishmeal as a source of DHA for n-3 PUFA enrichment of pork, chicken, and eggs. *Lipids*, 37: 1067-1076.
- Hooper, L., R. L. Thompson., R. A. Harrison., C. D. Summerbell., A. R. Ness., H. J. Moore., H. V. Worthington., P. N. Durrington., J. P. T. Higgins., N. E. Capps., R. A. Riemersma., S. B. J. Ebrahim and G. D. Smith (2006). Risks and benefits of omega 3 fats for mortality, cardiovascular disease, and cancer: systematic review. *BMJ Online First*, 1-9.
- Hugo, A., S. P. Els., F. H. de Witt., H. J. van der Merwe and M. D. Fair (2009a). Effect of dietary lipid sources on lipid oxidation of broiler meat. *South African Journal of Animal Science*, 39:149-152.
- Hugo, A., S. P. Els., C. Bothma., F. H. de Witt., H.J. van der Merwe and M. D. Fair (2009b). Influence of dietary lipid sources on sensory characteristics of broiler meat. *South African*
- Kris-Etherton, P.M., W. S. Harris and L. J. Appel (2002). Fish Consumption, Fish Oil, Omega-3 Fatty Acids, and Cardiovascular Disease. *Circulation*, 106:2747-2757.
- Kromhout, D., E. J. M. Feskens and C. H. Bowles (1995). The Protective Effect of a Small Amount of Fish on Coronary Heart Disease Mortality in an Elderly Population. *International Journal of Epidemiology*, 24(2): 340-345.
- Leskanich, C. and R. Noble (1997). Manipulation of the n-3 polyunsaturated fatty acid composition of avian eggs and meat. *World's Poultry Science Journal*, 53: 155-183.
- López-Ferrer, S., M. D. Baucells., A. C. Barroeta., A. Blanch and M. A. Grashorn (1999). N-3 enrichment of chicken meat using fish oil: Alternative substitution with rapeseed and linseed oils. *Poult. Sci.*, 78: 356-365.
- Lopez-Ferrer, S., M. D. Baucells., A. C. Barroeta and M. A. Grashorn (2001). n-3 Enrichment of Chicken Meat. 1. Use of Very Long-Chain Fatty Acids in Chicken Diets and Their Influence on Meat Quality: Fish Oil. *Poultry Science*, 80:741-752.
- Malayođlu, H.B., S. Özkan., S. Koçtürk., G. Oktay and M. Ergül (2009). Dietary vitamin E (α-tocopheryl acetate) and organic selenium supplementation: performance and antioxidant status of broilers fed n-3 PUFA-enriched feeds. *South African Journal of Animal Science*, 39:274-285.
- McMichael, A.J.M. and H. J. Bambrick (2005). Meat consumption trends and health: casting a wider risk assessment net. *Public Health Nutrition*, 8:341-343.
- Mellor, S. (2005). Omega 3 could enhance chicken meat. *WORLD POULTRY*, 21(7): 12-13.
- Mirghelenj, S.A., A. Golian and V. Taghizadeh (2009). Enrichment of Chicken Meat with Long Chain Omega-3 Fatty Acids though Dietary Fish Oil. *Research Journal of Biological Sciences*, 4(5): 604-608.
- Moghadasian, M.H. (2008). Advances in dietary enrichment with n-3 fatty acids. *Critical Reviews in Food Science and Nutrition*, 48: 402-410.

- Nieto, G. and G. Ros (2012). Modification of Fatty Acid Composition in Meat Through Diet: Effect on Lipid Peroxidation and Relationship to Nutritional Quality – A Review. In Angel Catala (Ed.), *Lipid Peroxidation* (pp. 239-258): InTech, DOI: 10.5772/51114.
- Pegg, R.B. and F. Shahidi (2012). Off Flavors and Rancidity in Foods. In Leo M. L. Nollet (Ed.), *Handbook of Meat, Poultry and Seafood Quality* (pp. 127-139): John Wiley & Sons Inc.
- Sacks, F. M. (2002). The role of high-density lipoprotein (HDL) cholesterol in the prevention and treatment of coronary heart disease. *Am. J. Cardiol.*, 15:139–143.
- Salma, U., A. G. Miah., T. Maki., M. Nishimura and H. Tsujii (2007). Effect of Dietary *Rhodobacter capsulatus* on Cholesterol Concentration and Fatty Acid Composition in Broiler Meat. *Poultry Science*, 86:1920–1926.
- Salamatdoustnobar, R., H. Aghdamshahriar and A. Gorbani (2008). Enrichment of Broiler Meat with n-3 Polyunsaturated Fatty Acids. *Asian Journal of Animal and Veterinary Advances*, 3:70-77.
- Sargent, J.R. (1997). Fish oils and human diet. *British Journal of Nutrition*, 78(1): S5-S13.
- Schacky, C.V. and W. S. Harris (2007). Cardiovascular benefits of omega-3 fatty acids. *Cardiovascular Research*, 73:310-315.
- Shahriar, H.A., A. Rezaei., A. Lak and A. Ahmadzadeh (2007). Effect of Dietary Fat Sources on Blood and Tissue Biochemical Factors of Broilers. *Journal of Animal and Veterinary Advances*, 6:1304-1307.
- Simopoulos, A.P. (1999). New Products from the Agri-Food Industry: The Return of n-3 Fatty Acids into the Food Supply. *Lipids*, 34:297-301.
- Smet, K., K. Raes., G. Huyghebaert., L. Haak., S. Arnouts and S. De Smet (2008). Lipid and Protein Oxidation of Broiler Meat as Influenced by Dietary Natural Antioxidant Supplementation. *Poultry Science*, 87:1682–1688.
- Speedy, A.W. (2003). Animal Source Foods to improve micronutrient nutrition in developing countries. *J. Nutr.*, 133: 4048S–4053S.
- Stone, N.J. (1996). Fish Consumption, Fish Oil, lipids, and coronary heart disease. *circulation*, 94: 2337-2340.
- Szymczyk, B., P. M. Pisulewski., W. Szczurek and P. Hanczakowski (2001). Effects of conjugated linoleic acid on growth performance, feed conversion efficiency, and subsequent carcass quality in broiler chickens. *British Journal of Nutrition*, 85:465-473.
- Tümová, E and Teimouri, A. (2010). Fat deposition in the broiler chicken : A review. *Scientia agriculturae bohemia*, 41(2):121- 128
- Walker, P., P. Rhubart-Berg., S. K. McKenzie and R. S. Lawrence (2005). Public health implications of meat production and consumption. *Public Health Nutrition*, 8:348–356.
- Zduńczyk, Z and J. Jankowski (2013). Poultry meat as functional food: modification of the fatty acid profile – a review. *Ann. Anim. Sci.*, 13(3): 463–480
- Zuidhof, M. J., M. Betti., D. R. Korver., F. I. L. Hernandez., B. L. Schneider., V. L. Carney and R. A. Renema (2009). Omega-3-enriched broiler meat: 1. Optimization of a production system. *Poult Sci.*, 88: 1108-1120.

## PhD Thesis Abstract

### Bio-control based management of brinjal shoot and fruit borer, *Leucinodes orbonalis* Guenee

V.Niranjana

Degree: Ph.D. in Agricultural Entomology

Chairman: Dr. R. Philip Sridhar

Institution: Tamil Nadu Agricultural University, Coimbatore

Year of Award: 2015

#### ABSTRACT

Brinjal shoot and fruit borer, *Leucinodes orbonalis* is the most serious pest of brinjal. As currently the farmers rely exclusively on the application of pesticides to control *L. orbonalis* the level of contamination exceeded the Maximum Residue Limit (MRL) in brinjal fruits in India. By considering these impacts of insecticides the present study was conducted to formulate bio-intensive pest management (BIPM) against *L. orbonalis*. The survey conducted at Coimbatore and Erode districts confirmed the availability of pupal parasitoid of *L. orbonalis*, *Trathalaflavo-orbitalis* in the fields. The study on seasonal incidence of *L. orbonalis* at Thondamuthur, Coimbatore during the year 2014 revealed that the maximum ( $r = -0.78^*$ ) and minimum ( $r = -0.43$ ) temperature negatively correlate with the damage intensity of *L. orbonalis* whereas Relative Humidity (RH) ( $r = 0.17$ ) and rainfall ( $r = 0.41$ ) positively correlate with the damage intensity. Further it was found that the maximum temperature played a major role in adult emergence and fluctuating the population of moth. The biology of *L. orbonalis* revealed that there was 5 larval instars in the lifecycle of *L. orbonalis* and the weather parameters existed in Rabi, 2013 ( $27 \pm 5^\circ\text{C}$ ) promoted the fecundity, incubation period and hatchability of eggs of *L. orbonalis* as well as developmental period and growth of all life stages of *L. orbonalis*. Screening of 35 brinjal germplasms showed that the germplasm with lower shoot thickness, short fruit pedicel and calyx with higher trichome density on lower surface of leaves were less susceptible against the infestation of *L. orbonalis* hence; priority should be given to these biophysical characteristics while developing host plant resistance varieties in breeding programmes. The pheromone trap efficiency studies revealed that Wota-T trap installed at crop canopy level with 3 mg lure concentration replaced at every 21 days were found effective in trapping the maximum number of *L. orbonalis* moths. Research on egg parasitoids of *L. orbonalis* is insufficient. Hence different *Trichogramma* species of egg parasitoids were tested for their efficacy against *L. orbonalis* under laboratory condition. Among the nine different species of *Trichogramma*, *T. pretiosum* and *T. embryophagum* performed well in laboratory and field conditions in reducing population of *L. orbonalis* with maximum yield in brinjal at field conditions. Further it was found that *T. pretiosum* (92%) and *T. embryophagum* (90%) had higher parasitization against the eggs of *L. orbonalis* whereas *T. chilonis*, *T. dendrolimi* and *T. evanescens* had no efficacy against *L. orbonalis*. Moreover, it was noted that the parasitoids preferred freshly laid eggs of *L. orbonalis* for parasitization and the selection of eggs for parasitization was reduced with the aging of host eggs. In addition, parasitoids showed various emergence rates and maximum emergence rate was observed in *T. embryophagum* (90.2%) and *T. pretiosum* (87.5%). Almost all *Trichogramma* sp. tested in this study were eco-friendly and protected the other natural enemies prevailed in brinjal cultivation. Results of pathogenicity test of three entomopathogenic fungi @  $2 \times 10^9$  conidia per ml and Bt k<sup>®</sup> showed maximum mortality by *Verticillium lecanii* against all larval instars of *L. orbonalis*. Among the plant product insecticides and FORS, NSKE 5% had higher efficacy in functioning as ovicide, larvicide and adulticide against the *L. orbonalis*. Acetamiprid 20% SP performed well among newer molecular insecticides tested against the eggs of *L. orbonalis*. It was noticed that almost all insecticides reduced the shoot infestation substantially in 14 days after 2<sup>nd</sup> spray. Chlorantraniliprole 18.5% SC was found to be the most effective insecticides in reducing shoot and fruit infestation in brinjal cultivation by *L. orbonalis*. Bio-Intensive Pest Management (BIPM) module with the management practices viz., installation of pheromone traps, seasonal release of *Trichogramma pretiosum* and *T. embryophagum*, spraying of *Verticillium lecanii*, NSKE 5%, Acetamiprid 20% SP and Chlorantraniliprole 18.5%, evaluated against *L. orbonalis* on brinjal recorded a lesser shoot infestation of *L. orbonalis* as compared to farmer's practice. BIPM was found to be superior to farmer's practice in all aspects in the brinjal field experiment.

**MPhil Thesis Abstract****Impacts of climatic parameters and crop management practices on induction of host plant resistance and population dynamics of beneficial insects towards management of viral diseases of chilli and tomato****K. Prasannath**

Degree: MPhil in Plant protection

Chairman: D. M. De Costa and K. S. Hemachandra

Institution: Postgraduate Institution of Agriculture, U.O.P

Year of Award: 2015

**ABSTRACT**

A study was conducted to determine the effects of climatic parameters and different crop management practices on induction of host plant resistance and population of beneficial insects towards management of viral diseases in chilli and tomato crops. Chilli (*Capsicum annuum* L.) variety MI Green and tomato (*Solanum lycopersicum* L.) variety Thilina were grown at five different locations of Sri Lanka, namely Kilinochchi, Mahailuppallama, Kundasale, Peradeniya and Rahangala under two crop management systems *viz.* pesticide-based management system and an IPM-based management system with less reliance on pesticides, during *maha* 2012/2013 and *yala* 2013 seasons. At each location, two crop management systems were tested in a nested treatment structure using a randomized complete block design with six replicates. Activity of defense enzymes, namely peroxidase,  $\beta$ -1,3-glucanase, chitinase, phenylalanine ammonia lyase and polyphenol oxidase was quantified spectrophotometrically in field grown chilli and tomato tissues collected at the first harvesting stage of the crops. Incidence of different types of virus diseases of tomato and leaf curl virus disease incidence of chilli were recorded over four crop growth stages, namely initial, crop development, mid season and late season. Abundance of different types of beneficial insects and herbivorous insects was quantified over the four different growth stages of two crops. Data on mean day temperature and daily rainfall were collected from each experimental site during the two cropping seasons. Relationships among virus disease incidence, defense enzymes activities of crops, population of beneficial insects and insect herbivores and climatic parameters were also analyzed. Findings of the study revealed that activity of all the defense enzymes in chilli and tomato tissues had a significant influence ( $p < 0.05$ ) by the cropping season, location, cropping season  $\times$  location interaction effect, location  $\times$  crop management practice interaction effect and cropping season  $\times$  location  $\times$  crop management system interaction effect. In general, there was no significant variation ( $p = 0.05$ ) between the two crop management systems on the activity of the enzymes in chilli and tomato tissues. It indicates the equal efficiency of both crop management systems on inducing the activity of enzymes in both types of crop tissues. There was no significant relationship ( $p = 0.05$ ) of virus disease incidence of the two crops with mean temperature of location or total rainfall of location. The beneficial insect population was significantly higher ( $p < 0.05$ ) in the two crops grown under IPM system than that under the pesticide management system. Moreover, the effect of management system on beneficial insect population was significantly influenced ( $p < 0.05$ ) by the location effect. A significant polynomial relationship ( $p < 0.05$ ) was observed between beneficial insect population and the mean temperature of location for both crops. No significant relationship ( $p = 0.05$ ) was found between the beneficial insect population and virus disease incidences of the two crops. Combined effect of virus disease incidence and herbivorous insect count had a significantly high influence ( $p < 0.05$ ) on the activity of defense enzymes in the two crop tissues. The IPM-based management system had lower environmental impact of pesticides compared to the pesticide-based management system used for both crops. Hence, it can be concluded that the IPM package tested in the study is as equally-capable as the pesticide-based management system in inducing defense-related enzymes in the plant tissues towards the management of virus diseases in chilli and tomato crops.

## PREPARATION OF MANUSCRIPTS

### Instruction to Contributors / Authors

#### General

Submission of manuscript to the Editor involves the assurance that it is original and that no similar paper, other than an abstract or a poster presentation, has been or will be submitted for publication elsewhere, without the consent of the Editorial Board. Research papers, read at Symposia and Reviews may be submitted. Research papers should describe original investigation or technological achievements. Reviews should be critical evaluations of existing knowledge in a specialized field. The journal also accepts research notes.

#### Presentation of Manuscripts

No maximum contribution is prescribed, but papers should be written clearly and concisely. All unnecessary textual matter, figures and tables must be eliminated. Manuscripts should be submitted in **Triplicate** including the original type written copy – type throughout in double spacing on one side of A4 paper with 4cm margin all sides and font size of 11pt. the typescript should be free of corrections. The papers should be divided in to section such as Abstract, Key words, Introduction, Material and Method, Result and Discussion, Conclusion, Acknowledgements and References.

#### Title and Author affiliation

The title should be concise and appropriately informative. The name of the institute where research was carried out should be stated with author's name. Present address of the authors should be given as footnotes.

#### Running Title

A short title not exceeding 45 characters should be provided for the running title.

#### Abstract

This should state concisely the scope of the work and the principal findings and should be suitable for direct use by abstracting journals. It should not exceed 200 words.

#### Key Words

A list of key words not exceeding five words choosen from the title, abstract or body of paper should be provided. They should be in alphabetical order and seprated by commas.

#### References

All references to publications made in the text should be presented in a list of references at the end of the paper. The manuscripts should be carefully checked to ensure that the spelling of author's names and dates are exactly the same in the text as in the reference list. In the text refer to the author's name (without initial) and year of publication. If reference is made in the text to publications written by more than two authors, the name of the first author should be used followed by *et al*. The list of references should be arranged alphabetically on authors' name in the list is also mentioned with co-authors, the following order should be used.

**Journal**

Arulnandhy, V., G.R. Bowers and O.D. Smith (1984) Evaluation of soybean genotypes for seed viability during storage. *Sri Lankan Journal of Agricultural Science* **21**:71-81. (Journal name should not be abbreviated)

**Book**

Little, T.H and F.J. Hills (1978) *Agricultural Experimentation*. John Wiley and Sons, New York, 350 pp.

**Sections of Book**

Gregg, B.R. (1982) Soybean seed quality and practical storage pp. 52 – 56. In: *Soybean seed quality and stand establishment* (Eds. J. B. Sinclair and J.A. Jacobs), INSTOY series NO. 22, University of Illinois, Illinois.

**Personal Communication**

In referring to a personal communication, the two words are followed by the year (eg. S. Sandanam, personal communication, 1993)

**Abbreviations and Symbols**

They are according to the recommendation in various parts of British standards, 1991: Letter symbol, sign and abbreviation should be used. Authors are encouraged to use SI system of units.

**Formulae and Equations**

Formulae should be type written, leaving ample space around the formulae. They should be started on the left margin and the number placed in parentheses to the right of the equation. Give the meaning of all symbols immediately after the equation in which they are first used.

**Research Notes**

The Journal may include a limited number of research notes. Authors should submit research notes only when they believe that rapid publication of their results is of utmost importance. A research note should not exceed 1,200 words, inclusive of illustration and tables. Research notes should be complete and suitable for citation with the title clearly indicating the contents.

**Submission of Manuscripts on CD**

The disk version may be submitted at the time of acceptance of the manuscript for publication with the revised manuscript, after minor revision. The disk copy and paper copy should match, i. e. the disk copy should be the version used to get the final print. The CD should be labeled with word processing package used, corresponding author's name and the name of the file.

**Copyright**

All papers once accepted and published in the journal become a copy of the Eastern University Sri Lanka. No part of the paper is permitted to be copied, edited or duplicated in any form without prior permission from the Editor/ AGRIEAST.

Manuscripts and all correspondence should be sent to:

**Editor, "AGRIEAST",**  
Faculty of Agriculture,  
Eastern University,  
Chenkalady – 30350,  
Sri Lanka.

Tel : +94652240530  
Fax : +94652240740  
E-Mail : agrieastusl@gmail.com  
Website : www. esn.ac.lk